



Exservan™

(riluzole) oral film

A different way to take riluzole

For people with ALS who have
difficulty swallowing some medications,
EXSERVAN™ may be an option.



RILUZOLE

REDESIGNED

*Medications are not actual size or depicted
in proportion to each other.*

Indication

EXSERVAN™ (riluzole) is indicated for the treatment of amyotrophic lateral sclerosis (ALS).

Important Safety Information

Do not use if you are allergic to riluzole or to any of its ingredients.

Please see full Important Safety Information on pages 10–11 and full Prescribing Information.

Swallowing difficulties may worsen over time



Difficulty swallowing, also referred to as dysphagia, is caused by the gradual weakening and eventual paralysis of the muscles in the face and throat.

Actor portrayal.

People with dysphagia sometimes crush tablets or take medication with food.

Generally, crushing tablets may result in the loss of medication



WHAT TO WATCH FOR

As ALS progresses, subtle changes can be signs that someone could be developing difficulty swallowing. Patients might learn to adapt, but **these are some signs to look for and talk to your care team about:**

- **Smaller bites or sips**
- **Shortness of breath while eating**
- **Longer mealtimes**
- **Frequent coughing or choking on food while swallowing**
- **A gurgling-sounding voice after swallowing**
- **Weight loss**

Swallowing difficulties in people with ALS may disrupt medication intake.

Taking riluzole with food can reduce how much of the medication is absorbed by the body. Riluzole should be taken at least 1 hour before or 2 hours after a meal



If your doctor determines it is appropriate, you may be prescribed EXSERVAN™

For people with ALS who have difficulty swallowing some medications, EXSERVAN™ provides:

- ✓ A similar medication as riluzole tablets
- ✓ A similar dosage strength as riluzole tablets
- ✓ A different way to take riluzole without having to swallow a tablet

Read the [Instructions for Use](#) in the FDA-approved patient labeling for complete directions on how to properly dose and administer EXSERVAN™ oral film.

Important Safety Information *(continued)*

Before using EXSERVAN™, tell your doctor about all the medicines you take and all your health conditions, including if you:

- Have liver problems.
- Are taking certain drugs known as strong or moderate CYP1A2 inhibitors such as ciprofloxacin, enoxacin, fluvoxamine, methoxsalen, mexiletine, oral contraceptives, thiabendazole, vemurafenib, zileuton.
- Are pregnant or intend to become pregnant during EXSERVAN™ therapy, or if you are breastfeeding or intend to breastfeed during EXSERVAN™ therapy.

Please see full Important Safety Information on pages 10-11 and [full Prescribing Information](#).



Exservan™ (riluzole) oral film

For people who have difficulty swallowing some medications, **EXSERVAN™ may be an option.**



*Not actual size.
For illustrative purposes only.*

Important Safety Information *(continued)*

EXSERVAN™ can cause serious side effects, including:

- **Liver Problems:** Cases of liver injury, some fatal, have occurred in patients taking riluzole. Consult your doctor promptly if you experience unexplained nausea, vomiting, stomach (abdominal) pain, fatigue, weight loss (anorexia), jaundice (yellowing of the skin or whites of the eyes), and/or dark urine.
- **Low White Blood Cell Counts (Neutropenia):** Tell your doctor if you develop a fever while taking EXSERVAN™.

An oral film formulation of riluzole that dissolves on the top of the tongue



- Rectangular in shape with a honey lemon flavor. Each film is individually packaged in a foil pouch
- Absorbed by the body at a similar rate as riluzole tablets and delivers a similar amount of medication
- EXSERVAN™ is riluzole
- Riluzole is widely prescribed for many people with ALS

Important Safety Information *(continued)*

- **Serious Lung Problems (Interstitial Lung Disease):** Tell your doctor if you have problems with your lungs or breathing such as dry cough and difficult or labored breathing. Discontinue EXSERVAN™ immediately if you develop interstitial lung disease.

Please see full Important Safety Information on pages 10-11 and full Prescribing Information.

EXSERVAN™ (riluzole) has the same dosing schedule as the tablet form of riluzole:
50 MG, TWICE A DAY.



Read the Instructions for Use in the FDA-approved patient labeling.

- EXSERVAN™ should be taken at least 1 hour before or 2 hours after a meal
- Do not cut or split the film
- Place the film on the top of the tongue, where it will stick and begin to dissolve (NOTE: Do not drink liquid, chew, spit, or talk while EXSERVAN™ is dissolving.)
- Average time to dissolve: 2 minutes
- Swallow saliva in a normal manner as EXSERVAN™ dissolves

Important Safety Information *(continued)*

The most common side effects include numbness in the mouth or tongue, muscle weakness, nausea, lung problems, high blood pressure, and stomach (abdominal) pain.

Exservan™ 
(riluzole) oral film

EXSERVAN™ (riluzole)
oral film has a
safety profile similar
to riluzole tablets

COMMON SIDE EFFECTS INCLUDE:

- Numbness in the mouth or tongue (38%)
- Muscle weakness (19%)
- Nausea (16%)
- Decreased lung function (10%)
- High blood pressure (5%)
- Abdominal pain (5%)



RILUZOLE PILL



EXSERVAN™ FILM

Not actual size. Size and shape of tablet may vary.

Important Safety Information *(continued)*


These are not all the possible side effects of EXSERVAN™. Consult your doctor for medical advice about side effects.

To report side effects, contact Mitsubishi Tanabe Pharma America, Inc. at 1-888-292-0058 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full Important Safety Information on pages 10-11 and full Prescribing Information.



*Not actual size.
For illustrative
purposes only.*



Important Safety Information *(continued)*

Use EXSERVAN™ as prescribed. The recommended dosage for EXSERVAN™ is 50 mg taken orally twice daily at least 1 hour before or 2 hours after a meal. Place EXSERVAN™ oral film strip on the top of the tongue where it will adhere and dissolve. Swallow in a normal manner. Do not cut or split the film or take liquids with EXSERVAN™. Do not chew, spit, or talk while EXSERVAN™ is dissolving. Read the Instructions for Use for EXSERVAN™.

Exservan™
(riluzole) oral film 



Important Safety Information

Indication

EXSERVAN™ (riluzole) is indicated for the treatment of amyotrophic lateral sclerosis (ALS).

Important Safety Information

Do not use if you are allergic to riluzole or to any of its ingredients.

Before using EXSERVAN™, tell your doctor about all the medicines you take and all your health conditions, including if you:

- Have liver problems.
- Are taking certain drugs known as strong or moderate CYP1A2 inhibitors such as ciprofloxacin, enoxacin, fluvoxamine, methoxsalen, mexiletine, oral contraceptives, thiabendazole, vemurafenib, zileuton.
- Are pregnant or intend to become pregnant during EXSERVAN™ therapy, or if you are breastfeeding or intend to breastfeed during EXSERVAN™ therapy.

EXSERVAN™ can cause serious side effects, including:

- **Liver problems:** Cases of liver injury, some fatal, have occurred in patients taking riluzole. Consult your doctor promptly if you experience unexplained nausea, vomiting, stomach (abdominal) pain, fatigue, weight loss (anorexia), jaundice (yellowing of the skin or whites of the eyes), and/or dark urine.
- **Low White Blood Cell Counts (Neutropenia):** Tell your doctor if you develop a fever while taking EXSERVAN™.
- **Serious Lung Problems (Interstitial Lung Disease):** Tell your doctor if you have problems with your lungs or breathing such as dry cough and difficult or labored breathing. Discontinue EXSERVAN™ immediately if you develop interstitial lung disease.



*Not actual size.
For illustrative purposes only.*

The most common side effects include numbness in the mouth or tongue, muscle weakness, nausea, lung problems, high blood pressure, and stomach (abdominal) pain.

These are not all the possible side effects of EXSERVAN™. Consult your doctor for medical advice about side effects.

To report side effects, contact Mitsubishi Tanabe Pharma America, Inc. at 1-888-292-0058 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Use EXSERVAN™ as prescribed. The recommended dosage for EXSERVAN™ is 50 mg taken orally twice daily at least 1 hour before or 2 hours after a meal. Place EXSERVAN™ oral film strip on the top of the tongue where it will adhere and dissolve. Swallow in a normal manner. Do not cut or split the film or take liquids with EXSERVAN™. Do not chew, spit, or talk while EXSERVAN™ is dissolving. Read the Instructions for Use for EXSERVAN™.

Please see full Prescribing Information including Patient Information at EXSERVAN.com.


Exservan™
(riluzole) oral film



Exservan™ (riluzole) oral film

The first and only oral film formulation of riluzole

EXSERVAN™ (riluzole) oral film is a different way to take riluzole. For people with ALS who have difficulty swallowing some medications, EXSERVAN™ may be an option.

Learn more about EXSERVAN™,
including how to access treatment, at

[EXSERVAN.com](https://www.exservan.com)

An ALS Education Specialist Is Ready to Help

We are ready to help you find the information you need to move forward on your journey with ALS and EXSERVAN™.

Call toll-free 1-833-630-1122 between 9 AM and 9 PM ET, Monday through Friday.

The ALS Education Specialist discusses basic information about Mitsubishi Tanabe Pharma America, Inc. products and does not take the place of your doctor. Be sure to talk to your doctor about all treatment-related questions, as your doctor is the best person to help you decide if treatment is right for you. If you have a medical emergency, please call 911. If you need to report an adverse event or product complaint, please call 1-888-292-0058.

Indication

EXSERVAN™ (riluzole) is indicated for the treatment of amyotrophic lateral sclerosis (ALS).

Please see full Important Safety Information on pages 10-11 and full Prescribing Information.

EXSERVAN and the EXSERVAN logo are trademarks of Aquestive Therapeutics, Inc. The corporate symbol of Mitsubishi Tanabe Pharma America is a registered trademark of Mitsubishi Tanabe Pharma Corporation.

All other company names, product names, trade/service marks or other trade names are the property of their respective owners.

For US audiences only.

© 2021 Mitsubishi Tanabe Pharma America, Inc. All rights reserved.

CP-EV-US-0102 08/21



Mitsubishi Tanabe Pharma America